

2019

Effect of a combined exercise program (aerobic and rebound therapy) with two different ratios on some physical and motor fitness indices in intellectually disabled girl

Amir Hossein Haghighi

Faculty of Sport Sciences, Hakim Sabzevari University, Sabzevar, Iran, ah.haghighi@hsu.ac.ir

Fatemeh Mohammadtaghipoor

Faculty of Sport Sciences, Hakim Sabzevari University, Sabzevar, Iran, ah.haghighi@hsu.ac.ir

Mohammadreza Hamedinia

Faculty of Sport Sciences, Hakim Sabzevari University, Sabzevar, Iran, ah.haghighi@hsu.ac.ir

Javad Harati

Faculty of Sport Sciences, Hakim Sabzevari University, Sabzevar, Iran, ah.haghighi@hsu.ac.ir

Follow this and additional works at: <https://www.balticsportscience.com/journal>



Part of the [Health and Physical Education Commons](#), [Sports Medicine Commons](#), [Sports Sciences Commons](#), and the [Sports Studies Commons](#)

Recommended Citation

Haghighi AH, Mohammadtaghipoor F, Hamedinia M, Harati J. Effect of a combined exercise program (aerobic and rebound therapy) with two different ratios on some physical and motor fitness indices in intellectually disabled girls. *Balt J Health Phys Act.* 2019;11(1):24-33. doi: 10.29359/BJHPA.11.1.03

This Article is brought to you for free and open access by Baltic Journal of Health and Physical Activity. It has been accepted for inclusion in Baltic Journal of Health and Physical Activity by an authorized editor of Baltic Journal of Health and Physical Activity.

Effect of a combined exercise program (aerobic and rebound therapy) with two different ratios on some physical and motor fitness indices in intellectually disabled girl

Abstract

Background: Individuals with intellectual disability (ID) are mostly neglected and hardly exercised. The purpose of this study was to investigate the effect of a combined program (aerobic and rebound therapy) with two different ratios on some physical and motor fitness indices in intellectually disabled girls. **Material and methods:** A total of 24 girls with ID (13–17 years old with an IQ of 50 to 70) were selected and randomly divided into two experimental and one control group. Experimental groups performed a combined aerobic rebound therapy program with two different proportions: (group 1 performed 10 minutes of aerobic exercises and 20 minutes of rebound therapy, group 2 performed 20 minutes of aerobic exercises and 10 minutes of rebound therapy) for 8 weeks (3 sessions per week). All physical and motor fitness tests including endurance performance, lower extremities endurance, cardio-respiratory endurance, static and dynamic balance, agility and coordination were taken before and after the training period. Data were analyzed using Shapiro-Wilk's tests, covariance and variance analysis with repeated measurements ($\alpha = 0.05$). **Results:** The results showed that both exercise programs in experimental groups improved endurance performance indices, lower extremities endurance, cardio-respiratory endurance, static and dynamic balance, agility and coordination in comparison to the control group. No significant differences were observed between the two experimental groups. **Conclusions:** In conclusion, to improve motor and physical fitness indices in girls with ID, both combination programs (aerobic-rebound therapy) with different proportions can be used.

Keywords

balance, exercise, fitness, health, intellectual disability

Creative Commons License



This work is licensed under a [Creative Commons Attribution-NonCommercial-No Derivative Works 4.0 License](https://creativecommons.org/licenses/by-nc-nd/4.0/).

Effect of a combined exercise program (aerobic and rebound therapy) with two different ratios on some physical and motor fitness indices in intellectually disabled girls

Authors' Contribution:

A Study Design
B Data Collection
C Statistical Analysis
D Data Interpretation
E Manuscript Preparation
F Literature Search
G Funds Collection

Amir Hossein Haghighi^{ACDEFG}, Fatemeh Mohammadtaghipoor^{BCEFG},
Mohammadreza Hamedinia^{ACDG}, Javad Harati^{CDEFG}

Faculty of Sport Sciences, Hakim Sabzevari University, Sabzevar, Iran

abstract

Background: Individuals with intellectual disability (ID) are mostly neglected and hardly exercised. The purpose of this study was to investigate the effect of a combined program (aerobic and rebound therapy) with two different ratios on some physical and motor fitness indices in intellectually disabled girls.

Material and methods: A total of 24 girls with ID (13–17 years old with an IQ of 50 to 70) were selected and randomly divided into two experimental and one control group. Experimental groups performed a combined aerobic-rebound therapy program with two different proportions: (group 1 performed 10 minutes of aerobic exercises and 20 minutes of rebound therapy, group 2 performed 20 minutes of aerobic exercises and 10 minutes of rebound therapy) for 8 weeks (3 sessions per week). All physical and motor fitness tests including endurance performance, lower extremities endurance, cardio-respiratory endurance, static and dynamic balance, agility and coordination were taken before and after the training period. Data were analyzed using Shapiro-Wilk's tests, covariance and variance analysis with repeated measurements ($\alpha = 0.05$).

Results: The results showed that both exercise programs in experimental groups improved endurance performance indices, lower extremities endurance, cardio-respiratory endurance, static and dynamic balance, agility and coordination in comparison to the control group. No significant differences were observed between the two experimental groups.

Conclusions: In conclusion, to improve motor and physical fitness indices in girls with ID, both combination programs (aerobic-rebound therapy) with different proportions can be used.

Key words: balance, exercise, fitness, health, intellectual disability.

article details

Article statistics: Word count: 3,539; Tables: 2; Figures: 1; References: 36

Received: August 2018; **Accepted:** November 2018; **Published:** March 2019

Full-text PDF: <http://www.balticsportscience.com>

Copyright © Gdansk University of Physical Education and Sport, Poland

Indexation: Celdes, Clarivate Analytics Emerging Sources Citation Index (ESCI), CNKI Scholar (China National Knowledge Infrastructure), CNPIEC, De Gruyter - IBR (International Bibliography of Reviews of Scholarly Literature in the Humanities and Social Sciences), De Gruyter - IBZ (International Bibliography of Periodical Literature in the Humanities and Social Sciences), DOAJ, EBSCO - Central & Eastern European Academic Source, EBSCO - SPORTDiscus, EBSCO Discovery Service, Google Scholar, Index Copernicus, J-Gate, Naviga (Softweco, Primo Central (ExLibris), ProQuest - Family Health, ProQuest - Health & Medical Complete, ProQuest - Illustrata: Health Sciences, ProQuest - Nursing & Allied Health Source, Summon (Serials Solutions/ProQuest, TDOne (TDNet), Ulrich's Periodicals Directory/ulrichsweb, WorldCat (OCLC)

Funding: This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Conflict of interests: Authors have declared that no competing interest exists.

Corresponding author: Amir Hossein Haghighi, Faculty of Sport Sciences, Hakim Sabzevari University, Sabzevar 96131, Iran; phone no.: +985144012765; E-mail: ah.haghighi@hsu.ac.ir.

Open Access License: This is an open access article distributed under the terms of the Creative Commons Attribution-Non-commercial 4.0 International (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non-commercial and is otherwise in compliance with the license.

INTRODUCTION

Intellectual disability is a complex dysfunction that occurs at different stages of development, during pregnancy, childbirth and postnatal, which can be due to diseases, infection and viruses, and it cannot be contagious [1]. This disorder, in addition to cognitive functions, disrupts motor activity of people with ID [2]. The functional disability of these individuals affects all aspects of their lives, such as behavioral, self-perceptual, and interpersonal relationships, and consequently it significantly reduces the quality of their life in society [2, 3]. Educable intellectually disabled children have a lower level of muscular strength, endurance, agility, running speed, reaction time and balance in comparison with their peer normal children. This imbalance of muscular strength and endurance changes the alignment of the body and increases the pressures and/or stresses on joints and tissues [4-6]. Golubovic et al. [6] reported that intellectually disabled people exhibited low performance in strength and muscular endurance, flexibility, motor coordination and cardiovascular endurance in standardized fitness tests. Therefore, performing physical activity plays an important role in improving the health indicators as well as reducing the specific health issues of intellectually disabled people. It has been emphasized that regular physical activity is an important component of a healthy lifestyle in these people [6, 7]. Moreover, researchers have argued that increased and sustained participation in moderate to vigorous physical activity can be one of the most effective ways to improve health in this population [5, 8]. However, the way of exercising to gain the most benefits and develop physical fitness and overall health depends on the proper manipulation of the volume, intensity and type of exercises. On the other hand, international and national organizations recommend that children with ID should be involved in physical activities. But there is some evidence of daily physical activity in these individuals and few specific standards and guidelines for them. According to the guidelines of the American College of Sports Medicine (ACSM) related to aerobic exercises for healthy individuals, it is necessary to perform a continuous physical activity with moderate intensity for 20-60 minutes and 3-5 times per week in order to keep cardiovascular fitness, body composition and muscle strength [9]. However, few studies have been conducted regarding other types of exercise programs and/or designing such programs for intellectually disabled individuals.

Rebound therapy is one of the enjoyable and attractive exercises for rehabilitation of people with ID. These exercises were first used to rehabilitate children with physical disabilities and learning difficulties [10]. Furthermore, it has been shown that in a steady state of the heart rate and oxygen uptake, more work can be done on a trampoline compared to treadmill and/or other types of physical activity, which means that oxygen uptake on a trampoline is more efficient [11]. Khalil Tahmasebi et al. [10] claimed that sequential trampoline jumps would reduce the risk of injury during landing. For this reason, this program could be considered as a suitable way to increase balance, muscle strength, vertical jump, and even treatment in injuries. On the other hand, aerobic exercises are defined as any kind of activity that requires the consumption of substantially more oxygen as well as a faster heart rate than at rest. Researchers have also paid special attention to the role of aerobic exercises in improving physical and cardiorespiratory fitness in people with ID [5, 13].

Considering the fact that exercises have improved physical fitness and function in intellectually disabled individuals, less attention has been paid to the

subject of how different training programs can lead to the best performance. In this regard, a combination of aerobic exercise and rebound therapy can be mentioned. Combining exercises may have synergistic effects of both types of exercise programs and can improve the performance of children with ID. However, what kind of combination of these exercises can provide more benefits for them is unclear and needs to be considered. Moreover, these children's motivations are low and they are less likely to participate in social and community programs, hence the implementation of a combination of exercises that creates diversity and joy which will help them to overcome their lack of motivation. Existing research has examined the effects of training programs individually and on the limited indicators of physical fitness in this population [2, 3, 14-17]. To the best of our knowledge, there is no study that has investigated combining these two programs. Therefore, the purpose of this study was to investigate the effect of a combined exercise program (aerobic and rebound therapy) with two different ratios on some physical and motor fitness indices in intellectually disabled girls.

MATERIAL AND METHODS

PARTICIPANTS

This research was a semi-experimental study. The statistical sample included 24 girls, from the Khamseh Gorjestani school in Quchan city, aged between 13-17 years old and with the IQ of 50 to 70. These individuals are targeted based on entry criteria including the absence of medical problems, the lack of multiple disabilities and specific diseases such as epilepsy, the absence of injury in lower extremities, and the ability to perform sports activities. Then subjects were randomly divided into three equal groups: 1) experimental group 1 (10 minutes aerobic and 20 minutes rebound therapy), 2) experimental group 2 (20 minutes aerobic and 10 minutes rebound therapy) and 3) control group. After completing the administrative requirements, obtaining permission and in coordination with the management of a special school in the Quchan city, with the help of the school counselor and according to the criteria of entry and exit, students were selected. After explaining the method and purpose of the research, the subjects' parents completed the medical history questionnaire and filled in the consent form. Before performing the exercise protocol, the variables of functional endurance, endurance of lower extremities, cardiorespiratory endurance, static and dynamic balance, agility, and neuromuscular coordination were taken by the researcher from all subjects. Then subjects of the experimental groups followed an 8-week course (3 sessions per week) of a combined exercise program. All measurements were repeated 48 hours after the last training session.

EXERCISE PROTOCOL

The training program was designed to motivate students to perform, and to form leisure activities and sports hours at school. The program included 15-10 minute warm-up including jogging and stretching exercises, then 30 minutes of training which for experimental group 1 included 20 minutes exercises on a trampoline and 10 minutes play, and vice versa for experimental group 2. Notably, training on the trampoline was done periodically in the form of one minute activity and one minute rest. Furthermore, 5 minutes were allowed for cooling down. One minute was added to the time of training to keep the principle of overload.

MEASURING THE RESEARCH INDICATORS

Endurance performance: To measure this index, 6 minute walk test was used [18].

Lower extremities endurance: Leg lift test was used to measure lower extremity endurance [4].

Cardiorespiratory endurance: The Rockport Fitness Walking Test was used to measure this variable [19].

Static balance: To measure this index, the subjects stood on a balance board with one leg (dominant leg) with closed eyes while holding their hands on the hip for 10 seconds, then the duration of their balance was recorded [10].

Dynamic balance: To measure this index, the subject walked on a balance board six steps heel to toe while their hands were on their hips, then the correct steps were calculated and recorded [10].

Agility: To measure agility, we used a test shown in Figure 1. In this test, a cone was located in the center and four were placed around it at three-meter distance. The subject moves from the center in the clockwise direction, after touching cone number 2, returns to the center, then goes to cone number 3. This process continues until the subjects goes to number 5 and passes the final line. Then the time is calculated [20].




Fig. 1. The test of agility [www.topendsports.com]

Coordination: Nelson's motion test was used. In this test, the two lines on each side were spaced 12 meters and 80 centimeters with a line in the middle. While standing the subject is prepared on the middle line facing the trainer. The examiner standing in front of the athlete and holding a timer in one hand shows one of the two directions with his other hand and immediately uses the stopwatch. The subject begins to run quickly to cross the finish line. At the same time, the examiner stops the time and records the athlete's result. This movement is repeated 6 times, 3 times randomly to the right and left. The interval between each repetition is 20 seconds. The average of 6 repetitions is considered as the final result of the subject [21].

STATISTICAL ANALYSIS

The Shapiro-Wilk test was used to determine the normal distribution of data, and descriptive statistics were used to display the central tendency and

dispersion. The ANOVA was used to examine the intragroup changes, and one-way ANCOVA and Bonferroni's post-hoc test was used to examine the changes between groups. All statistical operations were performed by SPSS software version 20, and the level of significance was set at $p \leq 0.05$.

RESULTS

Using the Shapiro-Wilk test showed that the distribution of all the variables is natural ($p < 0.05$). Therefore, parametric tests were used for statistical calculations. The characteristics of the subjects are presented in Table 1. This table shows that there is no meaningful difference between the groups in these indices. The results of statistical tests for the research variables are presented in Table 2.

Table 1. Characteristics of the subjects in the three groups

Exercise groups	P value	Control	Experimental 2	Experimental 1	P value
Height (m)	0.78	1.5 \pm 0.07	1.595 \pm 0.03	1.57 \pm 0.04	0.78
Weight (kg)	0.07	51.87 \pm 13.76	56.62 \pm 13.07	51.5 \pm 10.6	0.07
Body mass index (kg/m ²)	0.65	22.91 \pm 5.32	22.23 \pm 4.684	20.78 \pm 4.75	0.65

Significance level: $p \leq 0.05^*$

Table 2. Results of statistical tests on research variables

Variables	Groups	Pre-test	Post-test	Intragroup changes		Intergroup changes	
				F	P	F	P-value
Endurance performance (m)	Experimental 1	620 \pm 40.58	747.87 \pm 100.02	19.49	0.003*	16.06	0.001*
	Experimental 2	534.75 \pm 119.20	634.25 \pm 46.12	5.62	0.05*		
	Control	542.5 \pm 75.93	515.25 \pm 64.08	4.8	0.6		
Endurance of lower extremities (repetitions)	Experimental 1	6.75 \pm 3.45	11.25 \pm 5.49	29.84	0.001*	14.08	0.0001*
	Experimental 2	5.12 \pm 2.23	8.87 \pm 3.18	82.89	0.0001*		
	Control	3.12 \pm 2.41	2.87 \pm 2.23	0.63	0.45		
Cardio-respiratory endurance (ml/kg/min)	Experimental 1	29.72 \pm 0.28	30.29 \pm 0.43	8.44	0.023*	6.97	0.002*
	Experimental 2	29.7 \pm 0.58	29.95 \pm 0.64	1.35	0.28		
	Control	29.58 \pm 0.43	29.15 \pm 0.51	12.44	0.01*		
Dynamic balance (s)	Experimental 1	3.25 \pm 1.48	6.75 \pm 3.19	13.19	0.008*	6.44	0.007*
	Experimental 2	3.5 \pm 1.60	5.37 \pm 2.5	11.66	0.01*		
	Control	3.5 \pm 1.51	3.25 \pm 2.05	0.17	0.68		
Static balance with closed eyes (s)	Experimental 1	1.62 \pm 0.91	4 \pm 1.69	40.11	0.0001*	10.06	0.001*
	Experimental 2	1.87 \pm 0.83	3.12 \pm 0.83	9.21	0.019*		
	Control	1.5 \pm 1.06	1.62 \pm 1.3	0.3	0.59		
Agility (s)	Experimental 1	13.87 \pm 2.30	10 \pm 1.17	28.44	0.001*	18.7	0.0001*
	Experimental 2	16.15 \pm 4.23	11.6 \pm 1.8	12.87	0.009*		
	Control	15.59 \pm 4.10	16.21 \pm 4.02	7.5	0.02*		
Coordination (s)	Experimental 1	3.26 \pm 0.52	2.46 \pm 0.36	42.01	0.0001*	9.73	0.001*
	Experimental 2	3.55 \pm 0.93	2.95 \pm 0.36	7.5	0.02*		
	Control	3.89 \pm 0.30	3.99 \pm 0.9	0.11	0.78		

Significance level: $p \leq 0.05^*$

Table 2 shows that both experimental group 1 (rebound therapy – aerobic exercise) and experimental group 2 (aerobic exercise – rebound therapy) improved endurance performance indices, lower extremities endurance, cardio-respiratory endurance, dynamic balance, static balance with eyes closed, agility and coordination in comparison with the control group ($p \leq 0.05$). There was no significant difference between the two experimental groups ($p < 0.05$).

DISCUSSION

The present study showed that both combinations of exercises of rebound therapy and aerobic exercise significantly improved endurance performance and cardiorespiratory endurance indices in intellectually disabled girls compared with the control group. To our knowledge, no research on a combination of training programs of aerobic-rebound therapy on girls has been conducted so far. Therefore, we have used all the research that has been related, directly and indirectly, to the subject under discussion. With this explanation in the same field, Haghighi et al. showed that eight weeks of aerobic exercise, four times per week, significantly increased the traveled distance in a 300-yard walking test in girls with intellectual disability in comparison with exercises performed three times a week. They indicated that the reason was the increase in peak oxygen uptake (VO_{2peak}) and losing weight [22].

Wu et al. [14] concluded that a circular training program for 12 weeks, five times per week, improves cardiovascular fitness in obese children with ID. Oviedo et al. [23] demonstrated that a combined training program including aerobic, strength and balance exercises for 14 weeks improves cardiovascular fitness in people with ID. Furthermore, Boer et al. [16] stated that performing interval training for 15 weeks, twice a week, compared with continuous aerobic exercise, improves physical fitness of intellectually disabled teenagers. Other researchers also reported similar results in their studies on the same subject [4, 24, 25]. In addition, Ghasemi et al. [26] investigated the improvement in cardiorespiratory endurance after eight weeks of rebound therapy exercises in children with ID. Sadeghi et al. [11] reported that performing eight weeks of rebound therapy just as aerobic exercises could improve aerobic capacity in men with asthma [11]. Also, Cugusi et al. [27] showed that jumping on a trampoline by overweight women in one session can speed recovery in vigorous sports. In contrast, Nasrabadi et al. [28] showed that eight weeks of combined training (three times a week) had no significant effect on cardiorespiratory fitness and maximum oxygen uptake in intellectually disabled girls. They suggested the probable cause of this result would be insufficient training time. But, in the present study both training programs provided sufficient stimulus for the desired changes in the subjects' endurance and aerobic performance indices, since these two indicators can be influenced by both physiological and functional adaptations, such as increasing cardiac output, plasma volume, cardiac contractility, stroke volume, arterial-venous blood oxygen difference, muscle oxidative capacity, and muscle capillarity [29] and also include changes that occur in other physical and motor fitness components, such as static and dynamic balance, coordination, agility and muscular endurance which all in all can develop endurance and aerobic performance indirectly.

Moreover, the present study showed that both programs of combined exercise, predominantly rebound therapy and aerobic exercise, significantly improve the lower extremities muscular endurance in intellectually disabled girls in

comparison to the control group. Ahmadi and Daneshmandi [24] showed that a core stability training program for 6 weeks, three times a week, improves muscular endurance in boys with ID. Asgari [25] also stated that 10 weeks' water training improves lower body muscle endurance in boys with ID.

Golubovic et al. [6] showed that a six-month program three times per week for 45 minutes of moderate physical fitness could improve muscle strength and endurance in children with ID. Ghasemi et al. [26] also showed that a program of rebound therapy for eight weeks, three times a week, developed muscle endurance in 7-11-year-old children with ID. Others have shown improved muscular endurance by doing different exercises [4, 14]. The researchers attributed this finding to the relationship between increased strength and muscular endurance, since possibly strength training has increased muscle endurance to some extent, but a greater increase in muscular endurance depends on the specificity of training. In other words, muscle endurance can be increased by lowering exercise intensity and increasing the duration or repetitions of a single movement. Since the subjects in the present study performed 30 minutes continuous activity, improvement in muscle strength and endurance which has been observed during the period of the study is due to this matter. On the other hand, perhaps muscle recruitments and use of core muscles during running and trampoline movements, as well as the continuous use of lower limb muscles in daily activities have improved muscular endurance.

The other result of the present study was that both combination exercises with predominantly rebound therapy and aerobic exercise significantly improved the static and dynamic balance in comparison with the control group. Kachouri et al. [17] demonstrated that a combined strength and proprioceptive training program for eight weeks improved static balance with closed and opened eyes in boys with ID. Asgari [25] also showed an improvement in this indicator with 10 weeks' water training in mentally retarded boys. Wu et al. [14] observed an improvement in dynamic balance by implementing a 12-week circular exercise program in obese mentally retarded boys. Moreover, Khaliltahmasebi et al. [10] showed that doing eight weeks of rebound therapy exercises will improve the static and dynamic balance of educable children with intellectual disability. Giagazoglou et al. [30] also found that 12 weeks of trampoline training could be useful for developing the balance and motor function of children with mental disabilities. They have also shown that 12-week training on a trampoline improves balance in children with developmental coordination disorder [31]. Najafifard et al. [32] have shown that eight weeks of rebound therapy exercises improve the static and dynamic balance of male students with hearing impairment. They indicated this as a positive effect of muscle vibration that is caused by the use of trampolines, because the trampoline can create an unstable surface that causes more muscle recruitment and thus improves balance. Balance improvement in this study may be due to development in muscle strength and changes in the sensory-motor system of individuals participating in both training programs. In addition to the mentioned effect of various exercises, we can point out to neuromuscular adaptation, especially in the lower extremities, such as increasing the velocity of nerve conduction, increasing coordination between the agonist and antagonist muscles and reducing activity of the Golgi tendon organ [33].

This research showed that both programs of combined exercises, predominantly rebound therapy and aerobic, significantly improved the agility index compared

with the control group. Ghasemi et al. [26] showed that a rebound therapy program could improve agility in intellectually disabled children. Ilker et al. [4] showed that performing 10 weeks of swimming and water exercises improves the agility index in mentally retarded subjects [4]. Ahmadi and Daneshmandi [24] also found that six weeks of core stability training can improve agility in mentally retarded students. On the other hand, Faal Moghanlou et al. [34] did not report any significant effect on the agility index after 24 sessions of Spark motor program in the mentally retarded boys. The researchers highlighted the subjects' age, their low motivation and insufficient speed index as the explanation. The velocity of the stretch-shortening cycle is necessary for changing the speed and direction (agility). The rate of muscle contraction is recognized due to factors such as arousal and contractility of the musculoskeletal system, the number of active sarcomeres along the muscular fibers, the percentage of fast-twitch and slow-twitch fibers of the motor unit, and adenosine triphosphate (ATP) activity [35]. This mechanism is likely to improve agility performance in both groups of subjects. In addition, balance is one of the most important determinants of agility in men. In fact, agility consists of two components, speed and strength along with maintaining balance and coordination, thus, these indicators have developed in both training groups and improved the agility.

Another result of this study is that both combination programs of predominantly rebound therapy and aerobic exercise significantly improved the coordination index compared with the control group. Giagazoglou et al. [31] stated that 12-week training on a trampoline improves coordination in children with developed coordination disorders. Moghanlo et al. [34] showed that performing 24 sessions of the Spark motor program improves gross motor skills, such as double-sided coordination in educable mentally retarded boys. Asgari [25] also showed an improvement in the coordination index due to the implementation of 10 weeks of swimming and water exercises in boys with ID. Mitsiou et al. [36] stated that a training program with a trampoline improves neuromuscular coordination in 6-11-year-old students with developed coordination disorder. In contrast, Ghasemi Kahrizsangi et al. [33] showed that implementation of a rhythmic motor program in 8 weeks, three sessions per week, has no significant effect on perceptual-motor abilities of intellectually disabled girls. The researchers stated that the type and volume of the training program as their limitation caused subjects to fail to benefit from the class time in the same way as other girls, which probably led to non-significant results. However, various factors can affect the growth of basic motor skills, such as environment, geographical location and practice opportunity. Practice opportunity by using various facilities and equipment can provide suitable field to develop the perception of movement, strengthen muscles and develop basic motor skills. Sensory motor integration resulting from sequential jumps and improved balance ability, all of which can be gained from aerobic and rebound therapy, allows children to play their favorite games with their peers. Moreover, enhanced learning and experience of subjects in executing of various training programs as well as the development of other indicators affecting coordination, such as agility and balance in both training groups, has been effective in these results.

CONCLUSIONS

It is recommended that coaches use the two combinations of aerobic training and rebound therapy with different ratios to improve the quality of physical and motor fitness of intellectually disabled girls by taking into account their interests and training conditions.

This study did not distinguish different types of intellectual disability and possible differences between genders. Future studies can focus on this issue. Due to the fact that the length of the training program can affect the benefits gained by subjects, it is recommended that researchers utilize protocols of more than 30 minutes and 8 weeks in duration, then compare and evaluate the short- and long-term effects of the protocol.

AKNOWLEDGEMENTS

The authors wish to thank all participants who took part in the study.

REFERENCES

- [1] Afrooz GH. An introduction to the psychology and education of exceptional children. Tehran: Tehran University; 2006, 25-48. Persian.
- [2] Giagazoglou P, Arabatzi F, Dipla K, Liga M, Kellis E. Effect of a hippotherapy intervention program on static balance and strength in adolescents with intellectual disabilities. *Res Develop Disabil.* 2012;33(6):2265-2270. <http://doi.org/10.1016/j.ridd.2012.07.004>
- [3] Jankowicz-Szymanska A, Mikolajczyk E, Wojtanowski W. The effect of physical training on static balance in young people with intellectual disability. *Res Develop Disabil.* 2012;33(2):675-81. <http://doi.org/10.1016/j.ridd.2011.11.015>
- [4] Ilker Y, Nevin E, Ferman K, Bulent A, Erdal Z, Zafer C. The effects of water exercises and swimming on physical fitness of children with mental retardation. *J Hum Kinet.* 2009;21(9):105-111. <http://doi.org/10.2478/v10078-09-0013-6>
- [5] Shin IS, Park EY. Meta-analysis of the effect of exercise programs for individuals with intellectual disability. *Res Develop Disabil.* 2012; 33(6):1937-1947. <http://doi.org/10.1016/j.ridd.2012.05.019>
- [6] Golubovic S, Maksimovic J, Golubovic B, Glumbic N. Effects of exercise on physical fitness in children with intellectual disability. *Res Develop Disabil.* 2012;33(2):608-614. <http://doi.org/10.1016/j.ridd.2011.11.003>
- [7] Rosetty-Rodriguez M. Physical activity may promote health status individuals with Down syndrome. *Obesity.* 2006;15(2):67-71.
- [8] Jansen DE, Krol B, Groothoff JW, Post D. People with intellectual disability and their health problems: a review of comparative studies. *J Intellect Disab Res.* 2004;48(2):93-102. <http://doi.org/10.1111/j.1365-2788.2004.00483.x>
- [9] Pollock ML, Gaesser GA, Butcher ID, et al. The recommended quantity and quality of exercise for developing and maintaining cardiorespiratory and muscular fitness and flexibility in healthy adults. *Med Sci Sport Exerc.* 1998;30(6):975-991. <http://doi.org/10.1097/00005768-199806000-00032>
- [10] Khalil Tahmasebi R, Ghasemi Gh, Faramarzi S. The effects of rebound exercises on static and dynamic balance in educable children with mental retardation. *J Res Rehab Sci.* 2014;9(6):1050-1062. Persian.
- [11] Sadeghi M, Ghasemi GH, Zolaktav V, Miralaie A, Salehi M. The effects of rebound therapy and aerobic training on aerobic capacity, plasma endotheline level and the quality of life in male patients with asthma. *J Res Rehab Sci.* 2012;8(4):1-10. Persian.
- [12] Anderssen SA, Cooper AR, Riddoch C, Sardinha LB, Harro M, Brage S. Low cardiorespiratory fitness is a strong predictor for clustering of cardiovascular disease risk factors in children independent of country, age and sex. *Eur J Cardiovasc Prevent Rehab.* 2007;14(4):526-531. <http://doi.org/10.1097/HJR.0b013e328011efc1>
- [13] Oppewal A, Hilgenkamp TI, Van Wijck R, Evenhuis HM. Cardiorespiratory fitness in individuals with intellectual disabilities – A review. *Res Develop Disabil.* 2013;34(10):3301-3316. <http://doi.org/10.1016/j.ridd.2013.07.005>
- [14] Wu WL, Yang YF, Chu IH, Hsu HT, Tsai FH, Liang JM. Effectiveness of a cross-circuit exercise training program in improving the fitness of overweight or obese adolescents with intellectual disability enrolled in special education schools. *Res Develop Disabil.* 2017;31(60):83-95. <http://doi.org/10.1016/j.ridd.2016.11.005>
- [15] Pastula RM, Stopka CB, Delisle AT, Hass CJ. Effect of moderate-intensity exercise training on the cognitive function of young adults with intellectual disabilities. *J Strength Cond Res.* 2012;26(12):3441-3448. <http://doi.org/10.1519/JSC.0b013e318270fc83>
- [16] Boer PH, Meeus M, Terblanche E, et al. The influence of sprint interval training on body composition, physical and metabolic fitness in adolescents and young adults with intellectual disability: a randomized controlled trial. *Clin Rehab.* 2014;28(3):221-231. <http://doi.org/10.1177/0269215513498609>
- [17] Kachouri H, Borji R, Baccouch R, Laatar R, Rebai H, Sahli S. The effect of a combined strength and proprioceptive training on muscle strength and postural balance in boys with intellectual disability: An exploratory study. *Res Develop Disabil.* 2016;31(53):367-376. <http://doi.org/10.1016/j.ridd.2016.03.003>
- [18] Nasuti G, Stuart-Hill L, Temple VA. The six-minute walk test for adults with intellectual disability: a study of validity and reliability. *J Intellect Develop Disab.* 2013;38(1):31-38. <http://doi.org/10.3109/13668250.2012.748885>

- [19] Draheim CC, Laurie NE, McCubbin JA, Perkins JL. Validity of a modified aerobic fitness test for adults with mental retardation. *Med Sci Sport Exerc.* 1999;31(12):1849-1854. <http://doi.org/10.1097/00005768-199912000-00024>
- [20] Sopa Ioan S, Pomohaci M. Testing agility skill at basketball team (10-12 years old). *Sci Mov Health.* 2016;16(1):103-109.
- [21] Gharakhanlou R, Kordi MR, Gaeeni AA, Alizadeh MH, Vaezmousavi MK. Tests for assessing physical fitness, skill and psychological of elite athletes from different sports disciplines. 1st ed. Tehran: Asre Entezar Publication; 2012. P. 72. Persian.
- [22] Haghighi AH, Gelardi N, Hamedinia MR. The effect of two different volumes of aerobic exercise training on some risk factors of cardiovascular diseases in girls with mental retardation. *Sport Biosci.* 2015;7(2):241-259.
- [23] Oviedo GR, Guerra-Balic M, Baynard T, Javierre C. Effects of aerobic, resistance and balance training in adults with intellectual disabilities. *Res Develop Disabil.* 2014;35(11):2624-2634.
- [24] Ahmadi R, Daneshmandi H. The effect of core stabilization exercises on factors physical fitness to mental retardation. *Medicina Sportiva.* 2013;4(1):2058-2062. Persian.
- [25] Asgari H. The effect of an Aquatic training program with short and long sets on some factors of physical and motor fitnesses in individuals with mental retardation. [Thesis of Master of Science]. Sabzevar: Faculty of Sport Science. Hakim Sabzevari University; 2016. Persian.
- [26] Ghasemi GHA, Rahimi N, Khalil Tahmasebi R. The effects of rebound exercises on health-related physical fitness in educable children with mental retardation. *Sadra Med Sci J.* 2016;4(4):231-244. Persian.
- [27] Cugusi L, Manca A, Romita G, Bergamin M, Di Blasio A, Mercurio G. Exercise intensity and energy expenditure during a mini-trampoline rebounding exercise session in overweight women. *Sci Sport.* 2017;32(1):e23-e28. <http://doi.org/10.1016/j.scispo.2016.06.006>
- [28] Nasrabadi S. The effect of selected exercise training duration on some fitness factors in mentally retarded females. [Thesis of Master of Science]. Sabzevar: Faculty of Sport Science. Hakim Sabzevari University; 2014. Persian.
- [29] Ghalavand A, Shakeryan S, Nikbakht M, Mehdipour A, Monazamnezhad A, Delaramnasab M. Effects of aerobic training on cardiorespiratory factors in men with type 2 diabetes. *J Diabet Nursing.* 2014;2(2):8-17. <http://doi.org/10.1016/j.ridd.2013.05.034>
- [30] Giagazoglou P, Kokaridas D, Sidiropoulou M, Patsiaouras A, Karra C, Neofotistou K. Effects of a trampoline exercise intervention on motor performance and balance ability of children with intellectual disabilities. *Res Develop Disabil.* 2013;34(9):2701-2707. <http://doi.org/10.1016/j.ridd.2014.09.010>
- [31] Giagazoglou P, Sidiropoulou M, Mitsiou M, Arabatzi E, Kellis E. Can balance trampoline training promote motor coordination and balance performance in children with developmental coordination disorder? *Res Develop Disabil.* 2015;36:13-19. <http://doi.org/10.1016/j.ridd.2014.09.010>
- [32] NajafiFard A, Hemayat Talab R, sheikh M. The effect of eight-week period of rebound therapy exercises on the balance of male students with hearing disabilities. *Int J Sport Stud.* 2015;5(11):1206-1211.
- [33] Ghasemi Kahrizangi GHA, Salehi H, Heydari L. The effect of a rhythmic motor program on perceptual-motor abilities of educable mentally retarded children. *J Develop Motor Learning.* 2012;9:75-92. Persian.
- [34] Faal Moghanlo H, Hosseini F, Mikaili Manee F. Effect of Spark Motor Program on the development of gross motor skills in intellectually disabled educable boys. *J Birjand Uni Med Sci.* 2013;20(3):262-270. Persian.
- [35] Gaieeni AA, Rajabi H. Physical fitness. 4th ed. Tehran: SAMT Publication; 2011, 255. Persian.
- [36] Mitsiou M, Sidiropoulou M, Giagkazoglou P, Tsimaras V. Effect of trampoline based intervention program in static balance of children with developmental coordination disorder. *J Sport Med.* 2010; 45(2):125-143. <http://doi.org/10.1136/bjism.2010.081554.52>

Cite this article as:

Haghighi AH, Mohammadtaghipoor F, Hamedinia M, Harati J.
 Effect of a combined exercise program (aerobic and rebound therapy) with two different ratios on some physical and motor fitness indices in intellectually disabled girls.
 Balt J Health Phys Act. 2019;11(1):24-33.
 doi: 10.29359/BJHPA.11.1.03