Baltic Journal of Health and Physical Activity

Volume 3 | Issue 2

Article 3

2011

Differentiation of the Body Composition in the Polish National Team Pentathletes

Wladyslaw Jagiello Department of Martial Arts, Jedrzej Sniadecki Academy of Physical Education and Sport in Gdansk, Poland, wjagiello1@wp.pl

Roman M. Kalina Jerzy Kukuczka Academy of Physical Education in Katowice, Poland

Marina Jagiello Jedrzej Sniadecki Academy of Physical Education and Sport in Gdansk, Poland

Follow this and additional works at: https://www.balticsportscience.com/journal

Part of the Health and Physical Education Commons, Sports Medicine Commons, Sports Sciences Commons, and the Sports Studies Commons

Recommended Citation

Jagiello W, Kalina RM, Jagiello M. Differentiation of the Body Composition in the Polish National Team Pentathletes. Balt J Health Phys Act. 2011;3(2):105-111. doi: 10.2478/v10131-011-0010-8

This Article is brought to you for free and open access by Baltic Journal of Health and Physical Activity. It has been accepted for inclusion in Baltic Journal of Health and Physical Activity by an authorized editor of Baltic Journal of Health and Physical Activity.

Differentiation of the Body Composition in the Polish National Team Pentathletes

Abstract

Background: Contemporary anthropological research confirms the fact that body composition is one of the elements differentiating athletes from persons not practising sport systematically. The differences in body composition - in many cases even extreme - also regard athletes in the context of the sports discipline they practise. The aim of the study was to determine the properties of body composition of the Polish national team pentathletes in comparison to persons of the same population not practising competitive sport. Material/Methods: The research comprised members of the Polish national team in modern pentathlon (n=10). The subjects were 18-23 (20±1.5) years of age with their body mass from 64 to 80 kg (73.27±6.1) and body height of 172-188 cm (180.11±6.22). The athletes had had 6-13 years (10.38±2.07) of training experience. Students of Warsaw Technical University (Poland) constituted the reference group. Measurements of 20 basic somatic features were made. The following indices were calculated: leanness, Rohrer's, BMI, Manouvrier's and pelvic-shoulder. Body density, total body fat, active tissue, the general body composition profile and internal proportions of the body composition were determined. Results: An analysis of internal proportions of the factors of athletes' body composition revealed differences in particular groups of features. The athletes' overall body size results from greater than mean values of the body height and the average ones of stoutness and the fat content. Conclusions: The proportions of features within the analysed factors proved a significant advantage of the upper limb over the lower one and the significant predominance of the musculature of the forearm.

Keywords

body composition, pentathletes, national team

Creative Commons License



This work is licensed under a Creative Commons Attribution-Noncommercial-No Derivative Works 4.0 License.

	Differentiation of the Body Composition in the Polish National Team Pentathletes [*]						
	DOI: 10.2478/v10131-011-0010-8						
Authors' Contribution: A – Study Design B – Data Collection C – Statistical Analysis	Władysław Jagiełło ¹ ^(A, B, C, D, E, F, G) , Roman M. Kalina ² ^(C, D, E, F) , Marina Jagiełło ^{1 (C, D, E, F)}						
D – Data Interpretation E – Manuscript Preparation F – Literature Search G – Funds Collection	Poland ² Jerzy Kukuczka Academy of Physical Education in Katowice, Poland						
	Key words: body composition, pentathletes, national team						
	Abstract						
Background: Material/Methods:	Contemporary anthropological research confirms the fact that body composition is one of the elements differentiating athletes from persons not practising sport systematically. The differences in body composition – in many cases even extreme – also regard athletes in the context of the sports discipline they practise. The aim of the study was to determine the properties of body composition of the Polish national team pentathletes in comparison to persons of the same population not practising competitive sport. The research comprised members of the Polish national team in modern pentathlon (n=10). The subjects were 18-23 (20±1.5) years of age with their body mass from 64 to 80 kg (73.27±6.1) and body height of 172-188 cm (180.11±6.22). The athletes had had 6-13 years (10.38±2.07) of training experience. Students of Warsaw Technical University (Poland) constituted the reference group. Measurements of 20 basic somatic features were made. The following indices were calculated: leanness, Rohrer's, BMI, Manouvrier's and pelvic-shoulder. Body density, total body fat, active tissue, the general body composition profile and internal proportions of the body						
Results:	composition were determined. An analysis of internal proportions of the factors of athletes' body composition revealed differences in particular groups of features. The athletes' overall body size results from greater than mean values of the body height and the average ones of stoutness and the fat content						
Conclusions:	The proportions of features within the analysed factors proved a significant advantage of the upper limb over the lower one and the significant predominance of the musculature of the forearm.						
Word count: 2322							
Tables: 1 Figures: 2	Received: February 2011 Accepted: May 2011						
References: 25	Published: June 2011						
Address for correspondence							

Dr hab. prof. nadzw. Władysław Jagiełło Academy of Physical Education and Sport Dep. of Martial Arts, 80-336 Gdańsk, Poland, ul. K. Górskiego 1 Phone: +4858 554-74-50, e-mail: wjagiello1@wp.pl

Research conducted within the statutory research of Jędrzej Śniedecki Academy of Physical Education and Sport in Gdańsk (AWFiS).

Introduction

Contemporary anthropological research confirms the fact that body composition is one of the elements which differentiate athletes from persons not practising sport systematically [1, 2, 3]. The differences in body composition – in many cases even extreme – also regard athletes in the context of the sports discipline they practise [4, 5, 6].

With the increase in the training experience, hence the level of sports mastery, the decrease in differentiation of morphological features within the given discipline or event occurs. That is why research on athletes classified as the world elite in a particular sports discipline provides the most accurate information on the characteristic properties of their build [4, 7].

However, such properties are not easy to determine in all disciplines. This regards, among others, pentathlon, in which particular events substantially differ not only as far as the coordination structure is concerned but also with reference to the method of energy supply for the effort.

On the grounds of these general premises, the main aim of the study was an attempt to determine the properties of body composition of the Polish national team pentathletes in comparison to persons of the same population (reference group) not practising competitive sport and to answer the following questions:

- o Which somatic features determine the athletes' specific build?
- o Which internal proportions of the body composition are the most characteristic of them?

Material and methods

The research comprised members of the Polish national team in modern pentathlon (n=10). The subjects were 18-23 (20 ± 1.5) years old, with their body mass from 64 to 80 kg (73.27 ± 6.1) and body height 172-188 cm (180.11 ± 6.22). The athletes had had 6-13 years (10.38 ± 2.07) of training experience.

Anthropometric measurements were carried out in consistence with the generally adopted rules [8] using standard instruments. Furthermore, the following five indices were calculated: leanness, Rohrer's, BMI, Manouvrier's and pelvic-shoulder.

Total body fat in the percentage of body mass was calculated according to Brożek and Keys's equation [9] and body density on the basis of measurements of subcutaneous fat, with the application of Piechaczek's anticipating equation [10]. In total, 20 basic somatic features were measured. The general body composition profile of members of the national team was determined by means of the method of standardising features. Students of Warsaw Technical University constituted the reference group [11]. Moreover, the variation coefficient (V%) and the significance of differences were calculated by means of the Student test.

The assessment of internal proportions of the body composition was made through calculating Perkal's natural indices [12] with Milicerowa's modifications [13]. To do that the following were determined: composition factors (m), index of the total body size (M), internal proportions of the body composition (natural indices for each of the composition factors), evenness of the composition (the intrapersonal variability index), the code of internal proportions of the group (the point scale of Perkal's natural indices), internal proportions of the body composition within each of the factors.

Results

Mean values of the studied features in modern pentathletes prove that the slightest differentiation regards body density ($V_{\%}$ =0.50) and the leanness index ($V_{\%}$ =1.80). The greatest variability was determined in reference to fat content ($V_{\%}$ =12.04), the shank perimeter ($V_{\%}$ =9.47) and body mass ($V_{\%}$ =8.37) – Table 1.

The obtained results lay at the root of creating a profile of body composition of members of the Polish national modern pentathlon team (Fig.1). Their body build is characterised by a substantial perimeter of the forearm – they exceed their peers in this respect by 0.78 of the standardised Z value – and by the length of the upper limb (0.54 Z). Also worth attention is the knee width – here pentathletes are inferior to their peers by 0.42 Z and the shank perimeter (-0.41).

Facture	Pentathletes, n=10					Students, n=165		Test
reature	Х	SX	V%	min	max	X ₂	SX	Test t
Standing body height	180.11	6.22	3.45	171.50	187.70	179.360	6.190	0.370
Sitting body height	94.96	3.36	3.54	89.50	99.50	93.860	3.060	***4.618
Upper extremity length	80.19	2.94	3.66	76.20	84.00	78.300	3.510	***6.9167
Lower extremity length	85.15	4.29	5.04	77.30	91.50	85.500	4.100	-1.097
Shoulder width	40.39	2.05	5.08	37.50	44.50	40.670	1.590	**-2.262
Pelvis width	28.35	1.03	3.65	27.00	30.00	28.440	1.460	-0.792
Elbow width	6.89	0.55	8.02	6.00	7.80	6.980	0.340	***-3.400
Knee width	9.63	0.45	4.67	9.00	10.60	9.820	0.450	***-5.424
Forearm perimeter	27.42	1.25	4.56	26.00	30.00	26.020	1.800	***9.991
Shank perimeter	35.92	3.40	9.47	27.00	38.50	36.860	2.300	***-5.250
Body mass	73.20	6.12	8.37	64.00	80.00	72.110	8.960	1.563
Density	1.06	0.01	0.50	1.05	1.06	1.058	0.007	***-3.763
Fat, %	16.55	1.99	12.04	13.44	18.74	15.660	2.740	***4.173
Active tissue, %	83.45	1.99	2.39	81.26	86.56	84.340	2.740	***-4.173
Body area	1.92	0.11	5.77	1.77	2.05	1.902	0.125	*1.815
Leanness index	43.71	0.79	1.80	42.22	44.54	43.210	1.660	***3.841
Rohrer's index	1.25	0.07	5.53	1.18	1.39	1.250	0.150	0.213
BMI index	22.54	1.01	4.47	20.73	24.35	22.400	2.460	0.713
Manouvrier's index	89.72	4.54	5.06	78.72	95.31	91.093	2.820	***-6.232
Pelvic-shoulder index	70.31	3.43	4.88	65.06	75.20	69.929	2.400	*2.015

Tab. 1. Features of body composition in members of the Polish national team in modern pentathlon and students of Warsaw Technical University and the significance of differences

* p<0.05 ** p<0.01 *** p<0.001

The proportions of studied athletes' bodies determined by means of the indices clearly indicate their leptosomatic build (Rohrer's index), more masculine one (pelvic-shoulder index), with correct body mass (BMI) and an average length of lower extremities (Manouvrier's index).

The values of the factors of body composition confirm on a larger scale the observations of standardised values of individual features. As far as the overall body size is concerned, pentathletes are the most similar to the control group (M=0.02). The most distinguishing factor for pentathletes is the length factor (m_1 =0.23). The fat content factor and the stoutness factor amount to -0.09.

An analysis of mutual proportions between the pentathletes' body composition factors reveals a relatively proportional build – the intragroup variability index is 0.33. Features of length (0.32) are a characteristic element of such body composition. The contribution of stoutness and fat content features (-0.11) is the most proportional.

Using Perkal's natural indices point scale [22], the following code of the athletes' internal proportions was obtained: 5-4-4. This means that the overall body size (M) results from greater than mean values of body length and the average ones of stoutness and fat content.

The calculation of internal proportions of body composition within each of the factors provides very important information on the Polish modern pentathlon national team members' body build. Natural indices of the somatic features within the factors were calculated by subtracting the m factor value from the standardised features. The dominance of the upper extremity length (0.31) over the lower extremity length (-0.32) is manifested in features expressing the length index. The contribution of the features of standing and sitting body height to the length factor is even (Fig.2).

There is a marked dominance of the forearm musculature (0.87) in features reflecting the stoutness index. The knee width (-0.33) and the shank perimeter (-0.32) are the least proportionate to the overall value of the factor.



Fig. 1. The profile of body composition of members of the Polish national modern pentathlon team (n=10).



Fig. 2. Natural indices of the somatic features within factors among the Polish national modern pentathlon team (n=10).

Discussion

The properties of body composition of Polish representatives in modern pentathlon revealed in course of the study are a resultant of two processes. On the one hand, the process of sports selection of the most talented athletes; on the other hand, the effect of the adaptation of the organism to affecting it external factors. In this context, mean values of the height and the body mass of Polish representatives in pentathlon are interesting when compared to data from literature. The mean body height of members of the Polish national modern pentathlon team was 180 cm and mass 73 kg and did not statistically differ in this respect from the reference group (students of Warsaw Technical University). One should remember, however, that students of Warsaw Technical University are characterised by the highest indices of biological development among academic youth [11].

Greater differences occur in reference to men training other sports disciplines. For example, the mean body height of Polish representatives in tennis was 182 cm and their mass was 71 kg [14]. Slightly lower values were also achieved in the group of the best Polish judokas – 180 cm and 93 kg [15] and wrestlers – 174 cm and 78 kg [16]. Studying the first-league team members, Stawiarski [17] quotes the mean body height as 181.9 cm and the mass of 78.7 kg for volleyball players and 187.6 cm and 82.2 kg for basketball players. Piechaczek and Łaska-Mierzejewska [18] obtained the mean body height of 178.3 cm and body mass of 76.8 kg in the Polish national football team. Slightly lower values were also obtained in the group of the best Polish weightlifters – 166.9 cm and 73.9 kg [19] and hockey players – 174.5 cm and 73 kg [20].

On the example of combat sports extensive scientific research confirms the general trend in body composition development revealed in this study, which points to well-build competitors with athletic body composition with a more masculine build [6, 15, 21].

Competitors of the Polish national modern pentathlon team are characterised by a much greater value of the forearm perimeter and the upper extremity length in comparison to non-training persons. In the case of athletes, surprising is also a greater value of the fat content (0.33Z) than in non-training persons. The difference between these groups is statistically significant. Probably such body composition in pentathletes is connected with the nature of effort in this sports discipline. The dominance of technical events and the fact of using sports equipment (épée, pistol) may influence the athletes' somatic build.

Pentathletes are the most similar to the control group in the overall body size. Still, the greatest advantage of the athletes is manifested in the length factor. Studies carried out on a group of national tennis team representatives [22] showed that the slightest differences in body composition (in comparison to students of Warsaw Technical University) regarded the fat content and the overall body size factors. The length factor proved to be the most distinguishing feature of the tennis players. The results of studies on Polish representatives in judo and wrestling [15, 16] require a more complex comparative analysis. Taking the specificity of these sports disciplines into consideration (competing in strictly specified weight categories) one can conclude that Polish modern pentathlon representatives are the most similar in their body composition to judokas and wrestlers from mid-weight categories. In the other categories the differences are significant and concern all composition factors.

Natural indices of body composition factors of Polish national team competitors in modern pentathlon indicate their harmonious build – the code of internal proportions is 5-4-4. This is also confirmed by a low value of the intrapersonal variability index. With regard to the harmoniousness of the composition, pentathletes exceed both tennis players [23] and Polish national judo [15] and wrestling [16] representatives. The results of studies on pentathletes do not confirm, however, the tendencies revealed in Łaska-Mierzejewska's studies [24]. On the example of sports games she found a tendency in highly-qualified female athletes to be characterised by a significant predominance of one factor over the other ones. National team representatives manifested very high values of the intergroup variability, while the third league athletes, especially volleyball players, had their indices close to zero. Top-class athletes' differentiation of body composition was also found by Claessens et al. [25].

Natural indices of the somatic features within the factors provide very important information on the specificity of body composition in athletes practising modern pentathlon. A marked dominance of the musculature of the forearm is the characteristic feature in the stoutness factor.

The dominance of the upper extremity length over the lower extremity one is apparent among features expressing the length factor. Similar results were obtained on the tennis material [22]. On the one hand, in both these disciplines athletes use sports equipment (a racket in tennis and an épée and a pistol in pentathlon) requiring precision and swiftness of action, which is connected with a firm and secure grip. On the other hand, athletes with distinctly longer upper limbs have a greater reach, which gives them an advantage during a fight.

The results of our studies have proved that, as far as their body composition is concerned, men with successes in modern pentathlon significantly differ from their peers not practising sports (reference to students of Warsaw Technical University) and from athletes practising other sports disciplines.

Conclusions

- 1. Long-term training influence as well as the selection process of the most predisposed athletes have a great impact on the somatic build of the representatives of the Polish national modern pentathlon team. A larger forearm perimeter and the upper extremity length are characteristic features of their body composition. Slightly higher fat content is also noteworthy.
- 2. Body proportions of the studied athletes determined by means of indices clearly indicate their build, which is leptosomatic (Rohrer's), more masculine (pelvic-shoulder), with correct body mass (BMI) and average lengths of lower extremities (Manouvrier's).
- 3. An analysis of internal proportions of factors of the body composition in the representatives of the Polish national modern pentathlon team revealed differences in particular groups. The athletes' overall body size results from a more than average intensity of length features and the average ones of stoutness and fat content (5-4-4).
- 4. The proportions of somatic features within the length factor show a significant advantage of the upper limb length over the lower one. The input of height features while standing and sitting into the length factor is equal.
- 5. Within features reflecting the stoutness factor, there is a marked dominance of the forearm musculature. The least proportional are the knee width (-0,33) and the shank perimeter (-0,32). The contribution of the remaining features of the factor is relatively proportional.

References

- 1. Carter JL. The somatotypes of olympic athletes. Presented paper; 1978.
- 2. Charzewski J, editor. Antropologia [in Polish] [Anthropology]. Warszawa: AWF; 1999.
- 3. Garay AL, Levine L, Carter JE. Genetic and anthropological studies of olympic athletes. New York San Francisco London: Academic Press; 1974.
- 4. Baxter-Jones ADG, Helms P, Maffulli N, Baines-Preece JC. Growth and development of male gymnasts, swimmers, soccer and tennis players: a longitudinal study. *Annals of Human Biology* 1995;5:381-394.
- 5. Tumanian GC, Martirosov EG. Teloslozhenye i sport [in Russian]. Moskva: Fizkultura i Sport; 1976.
- 6. Almansba R, Sterkowicz S, Belkacem R, Sterkowicz-Przybycień K, Mahdad D. Anthropometric and physiological profiles of the Algerian Olympic judoists. *Arch Budo* 2010;6(4):85-193.
- 7. Leake CN, Carter JEL. Comparison of body composition and somatotype of trained female triathletes. *J Sport Sci* 1991;9(2):125-135.
- Drozdowski Z. Antropologia sportowa Morfologiczne podstawy wychowania fizycznego i sportu [in Polish] [Sports anthropology. Morphological bases of physical education and sport]. Warszawa – Poznań: PWN; 1979.
- 9. Brożek J, Keys A. The evaluation of leanness fatness in man. Norm and interrelationships. *Brit J Nutr* 1949;5:194-206.
- 10. Piechaczek H. Oznaczanie całkowitego tłuszczu ciała metodami densytometryczną i antropometryczną [in Polish] [Marking total body mass by the densitometric and anthropometric methods]. *Materiały i Prace Antropologiczne* 1975;89:3-48.
- 11. Piechaczek H, Lewandowska J, Orlicz B. Zmiany budowy ciała młodzieży akademickiej Politechniki Warszawskiej w okresie 35 lat [in Polish] [Changes in body composition among academic youth at Warsaw Technical University during 35 years]. *Wychowanie Fizyczne i Sport* 1996;3:3-14.

- 12. Perkal J. O wskaźnikach antropologicznych [in Polish] [On anthropological indices]. *Przegląd Antropologiczny* 1953;19:209-211.
- 13. Milicerowa H. Zastosowanie wskaźników Perkala do charakterystyki budowy ciała bokserów [in Polish] [The application of Perkal's indices to characterise boxers' body composition]. *Materiały i Prace Antropologiczne* 1956;20.
- 14. Jagiełło W, Jagiełło M. The profile of the body type for Poland's male tennis representatives. In: Ermakov SS, editor. Physical Education of the Students of Creative Profession. Kharkiv: Kharvoskaya Gossudarstvennaya Akademya Dizayna i Isskustv; 2005, No7, 60-66.
- Jagiełło W, Kalina RM, Tkaczuk W. Morphological differentiation of judo competitors. In: Szopa J, Gabryś T, editors. Sport training in interdisciplinary scientific researches. Chapter II. Czestochowa: Faculty of Management Technical University of Czestochowa; 2004, 200-206.
- 16. Jagiełło W, Kruszewski A. Morphological diversification of competitors training Greco-Roman style of wrestling. *Arch Budo* 2009;5:147-153.
- 17. Stawiarski W. Wyniki sportowe a morfologia, wiek i staż zawodniczy w wybranych grach zespołowych [in Polish] [Sports Results and the Morphology, Age and Competition Experience in Selected Team Sports]. *Rocznik Naukowy AWF Kraków* 1975, t. XIII.
- 18. Piechaczek H, Łaska-Mierzejewska T. Zróżnicowanie morfologiczne zawodników piłki nożnej [in Polish] [Morphological Differentiation of Football Players]. *Wychowanie Fizyczne i Sport* 1977;3:113-123.
- Drozdowski Z. Antropologia sportowa: morfologiczne podstawy wychowania fizycznego i sportu [in Polish] [Sports anthropology: morphological bases of physical education and sport]. 3rd ed. Warszawa – Poznań: PWN; 1984.
- 20. Gowarzewski Z. Wybrane cechy somatyczne polskich hokeistów na tle czołówki swiatowej [in Polish] [Selected somatic features of Polish hockey players in comparison to the world leaders]. *Sport Wyczynowy* 1976;2:37-41.
- 21. Kuźmicki S, Jagiełło W. Niektóre różnice i podobieństwa w budowie ciała judoków [in Polish] [Selected similarities and differences in judoists' body composition]. in: Kalina RM, Jagiełło W, editors. Wychowawcze i utylitarne aspekty sportów walki [Educational and utilitarian aspects of combat sports]. Warszawa: AWF; 2000, 136-143.
- 22. Jagiełło M. The profile of the body type for Poland's female tennis representatives. In: Ermakov SS, editor. *Pedagogics, Psychology, Medical-Biological Problems of Physical Training and Sports*. Kharkiv Kharvoskaya Gossudarstvennaya Akademya Dizayna i Isskustv; 2005, № 23, 135-139.
- 23. Jagiello M, Jagiello W. Internal proportions of the body composition in members of the female national tennis team of Poland. *Education. Physical Training. Sport.* [Lithuanian Academy of Physical Education. Kaunas] 2009;2(73):28-34.
- 24. Łaska-Mierzejewska T. Antropologia w sporcie i wychowaniu fizycznym [in Polish] [Anthropology in sports and physical education]. Warszawa: Biblioteka Trenera. Centralny Ośrodek Sportu; 1999.
- 25. Claessens A, Buenen G, Lefevre J, Martens G, Wellens R. Body structure, somatotype and motor fitness of top-class Belgian judoists and karateka: a comparative study. In: Reilly T, Watkins J, Borms J, editors. *Kinanthropometry* III. London: Spon; 1986, 53.